

Index

- ABC approach 16
- adolescents 12, 14–15
- aggressive behaviour
 - behavioural strategies 44–49
 - definition and diagnosis 42
 - epidemiology 43
 - groups at risk 43
 - pharmacotherapy 24, 49–54
 - risks to self and others 44
 - staff and parent training 48–49
 - treatment interventions 42–56
- anticonvulsants 24, 51
- antidepressants 15, 23, 51–52
- antisocial behaviour 13
- anxiety 14, 23, 30, 35–36, 52
- anxiolytic and sedative drugs 52
- applied behaviour analysis (ABA)
 - 45–47
- Asperger's disorder 5, 6, 25–41
 - age of onset 26–27
 - attention disorders 40
 - case examples 34, 37–38, 39–40
 - causation 29
 - clinical features 29, 34–41
 - diagnosis 27–29, 32
 - differences from autism 30–31
 - dysexecutive syndrome 39–40
 - epidemiology 26
 - Fifth Child form 38, 40
 - intersubjectivity 38
 - lack of empathy 36–37
 - learning and memory disorders 40–41
 - motor functioning 31, 35–36
 - non-verbal expression 34–35
 - non-verbal interpretation 36–38
 - semantic-pragmatic disorder 38–39
 - symbolic and language disorders 41
 - unusual interests 35–36, 41
- assessment 6–8, 29
- attention deficit hyperactivity disorder 15, 24, 40
- attention disorders 40
- atypical neuroleptics 53
- augmentative communication 20–22
- autism 1–24
 - aggression and intellectual disability 43
 - assessment 6–8
 - associated features 11
 - associated medical conditions 4
 - brain research 2–3

- causation 1–4
- cognitive abilities and IQ 11–12
- communication skills 9–10
- course and outcome 12–13
- diagnosis 5–6
- differences from Asperger's disorder 30–31
- emotional and behavioural problems 13–15
- epidemiology 4–5
- genetic research 3–4
- high functioning 11–12, 26, 28–29, 30
- historical views 2
- play and imagination 10
- ritualistic and stereotyped interests/behaviours 10–11
- social impairments 8–9
- treatment approaches 15–24
- autism spectrum disorders 5–6
- behavioural approaches 15–19, 44–49
- behavioural problems 30
 - in autism 13–14, 18–19
 - communicative function 18–19
 - environmental minimisation 57–64
 - medication 24, 49–54
 - treatment for aggression 42–56
- benzodiazepines 52
- beta-blockers 52
- body language of carers/educators 63
- bupirone 23, 24, 52
- care environment 57–64
- carers/educators 48–49, 62–64
- causation 1–4, 29
- chaining 18
- classical conditioning 46
- clonidine 52
- clumsiness 31, 36
- cognitive deficits 11–12
- cognitive therapies 55
- communication impairments 9–10, 34–35, 36–38
- communication skills programmes 20–22
- communicative function of difficult behaviour 18–19
- Compic (computer-generated pictures) 21
- counselling 55
- cycling mood disorder 24
- daily programme 62
- depression 12, 14–15, 23
- diagnosis 5–6, 27–29, 32, 50–51
- disruptive behaviour 14, 18, 24, 30
- drugs 14, 15, 23–24, 49–54, 59
- dyscalculia (arithmetic impairment) 41
- dysexecutive syndrome 37, 39–40
- echolalia 9
- educational programmes 19–20
- education environment 57–64
- environment
 - behaviour of carers/educators 62–64
 - daily programme 62
 - importance 57–59
 - minimisation of disturbed behaviours 57–64
 - physical 61–62
 - possible improvements 59–64
 - social interaction 60–61

- epidemiology 4–5, 26
- epilepsy 4, 11, 12, 43
- executive functioning 30, 39–40

- fenfluramine 53
- Fifth Child syndrome 38, 40
- functional assessment 47–48
- functional communication training 47

- gender differences 3, 26
- genetic factors 2, 3–4, 29

- high functioning autism 11–12, 26, 28–29, 30
- instruments for assessment 8
- intellectual disability 11–12, 42–56
- interests 10–11, 35–36, 41
- intersubjectivity 38
- IQ scores 11–12
- islets of ability 11

- language impairment 9–10, 29, 38–39, 41
- lithium 24, 52

- Makaton 20
- medical conditions associated with autism 4
- medication 14, 15, 23–24, 49–54, 59
- memory disorders 40–41
- motor functioning 31, 35–36

- naltrexone 24
- nature theory 2
- neuroleptics 23, 24, 51, 53
- noise levels, environmental 61–62
- non-verbal communication 20–22, 34–35, 36–38, 39

- object timetables 22
- obsessive–compulsive symptoms 10–11, 23, 36
- operant conditioning 46–47
- opiate blockers 53

- perceptual abnormalities 11
- pervasive developmental disorders (PDDs) 5, 27, 28–29
- pharmacotherapy 59
 - for aggressive behaviour 49–54
 - in autism 14, 15, 23–24
- photographs 21
- physical environment 61–62
- picture scripts 22
- play and imagination 10
- positive reinforcement 16
- prompting 17
- psychiatric diagnosis 50–51
- psychogenic theory 2, 4
- psychological treatments 14, 15–19, 44–49, 55
- psychotherapy 55

- receptive language 9
- restraint 49
- ritualistic interests/behaviours 10–11, 35–36

- secretin 54
- selective serotonin reuptake inhibitors (SSRIs) 23
- semantic-pragmatic disorder 38–39
- Sensori-integration therapies (SIT) 55
- serotonin 3
- shaping 17–18, 46–47
- signing 20–21

Index

- social environment 60–61
- social impairments 8–9, 28
- social skills groups 22–23
- social skills programmes 22–23, 41
- social stories 22
- special educational programmes 19–20
- speech abnormalities 9, 29, 38–39
- stereotyped interests/behaviours 10–11, 35–36
- stimulants 24, 52
- theory of mind 38, 39
- time out 16–17
- treatment interventions
 - for aggressive behaviour 42–56
 - in autism 15–24
 - early intervention 19, 49
 - points to remember 24
- triad of social impairments 28
- tricyclic antidepressants (TCAs) 23, 24